

Appendix A

Proposed Section X

Of the number reported in Section D, Line 13 how many pregnancies resulted in maternal, infant or fetal deaths? Include terminations prior to 20 weeks gestation (miscarriages, abortions and abortions for medical reasons). Include deaths discovered **BOTH** at home and after transfer to another provider or facility. You will be given an opportunity to further elaborate on these deaths in future screens and details for each death will be entered separately.

Fetal deaths (pregnancy losses prior to birth of a live neonate)

1. Was the death discovered OOH? (If yes continue, If no skip to #2 below)
 - A. Which county was the death discovered in?
 - B. Was the death prior to 20 weeks gestation (if yes continue, if no skip to C below)
 - i. Was the loss the result of a fetal anomaly? (yes, no, unkn) *it is not required by statute to collect cause of death for fetal deaths prior to 20 completed weeks gestation*
 - C. Was the loss greater than 20 0/7 weeks gestation **AND** diagnosed by a physician **AND** by maternal request, subsequently delivered OOH by you? (if yes continue, if no skip to D below)
 - i. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (If yes continue, if no skip to (ii) below)
 - a. Was the loss prior to onset of labor or after rupture of membranes without labor, **AND** between 20 0/7 weeks gestation up to and including 36 6/7 weeks gestation **OR**
 - b. Was the loss prior to onset of labor or after rupture of membranes without labor **AND** after 37 0/7 weeks gestation **OR**
 - c. Was the loss during labor **AND** between 20 0/7 weeks gestation and 36 6/7 weeks gestation **OR**
 - d. Was the loss during labor **AND** after 37 0/7 weeks gestation
 - ii. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (no)
 - a. Was the loss prior to onset of labor or after rupture of membranes without labor, **AND** between 20 0/7 weeks gestation up to and including 36 6/7 weeks gestation **OR**
 - b. Was the loss prior to onset of labor or after rupture of membranes without labor **AND** after 37 0/7 weeks gestation **OR**
 - c. Was the loss during labor **AND** between 20 0/7 weeks gestation and 36 6/7 weeks gestation **OR**
 - d. Was the loss during labor **AND** after 37 0/7 weeks gestation
 - D. Was the loss greater than 20 0/7 weeks gestation **AND** diagnosed by a physician **AND** by maternal request, subsequently delivered OOH by you? (no)
 - i. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (no)
 - a. Was the loss prior to onset of labor or after rupture of membranes without labor, **AND** between 20 0/7 weeks gestation up to and including 36 6/7 weeks gestation **OR**

- b. Was the loss prior to onset of labor or after rupture of membranes without labor **AND** after 37 0/7 weeks gestation **OR**
 - c. Was the loss during labor **AND** between 20 0/7 weeks gestation and 36 6/7 weeks gestation **OR**
 - d. Was the loss during labor **AND** after 37 0/7 weeks gestation
- 2. Did the death occur after transfer to another provider or facility? (if yes continue if no return to (1) above)
 - i. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (If yes continue, if no skip to (ii) below)
 - a. Was the loss prior to onset of labor or after rupture of membranes without labor, **AND** between 20 0/7 weeks gestation up to and including 36 6/7 weeks gestation **OR**
 - b. Was the loss prior to onset of labor or after rupture of membranes without labor **AND** after 37 0/7 weeks gestation **OR**
 - c. Was the loss during labor **AND** between 20 0/7 weeks gestation and 36 6/7 weeks gestation **OR**
 - d. Was the loss during labor **AND** after 37 0/7 weeks gestation
 - ii. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (no)
 - a. Was the loss prior to onset of labor or after rupture of membranes without labor, **AND** between 20 0/7 weeks gestation up to and including 36 6/7 weeks gestation *we could include a question asking the exact number of gestational weeks* **OR**
 - b. Was the loss prior to onset of labor or after rupture of membranes without labor **AND** after 37 0/7 weeks gestation **OR**
 - c. Was the loss during labor **AND** between 20 0/7 weeks gestation and 36 6/7 weeks gestation *we could include a question asking the exact number of gestational weeks* **OR**
 - d. Was the loss during labor **AND** after 37 0/7 weeks gestation *we could include a question asking the exact number of gestational weeks*
 - e. Complication contributing to or causing death, mark all that apply. (this has not been collected prior to this but we believe it should be collected)
 - i. Infection
 - ii. Meconium aspirations, other respiratory
 - iii. Neurological issues/seizures
 - iv. As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - v. As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - vi. Other medical issue
 - vii. Unknown
 - viii. Information not obtainable
 - ix. Other

Neonatal deaths (live born infant who died prior to the 7th day of life)

1. Did the death occur OOH (if yes continue, if no skip to (2) below)

A. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if yes continue, if no skip to (b) below)

a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*

Complication contributing to or causing death, mark all that apply

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

B. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if no continue, if yes revert to (A) above)

a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries

- As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other
- b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)
Complication contributing to or causing death, choose **ONE** from the following list
- Infection
 - Meconium aspiration, other respiratory
 - Neurological issues/seizures
 - As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other
2. Did the death occur after transfer to another provider or facility? (if yes continue, if no revert to (1) above)
- A. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if yes continue, if no skip to (b) below)
- a. Was neonate born of a gestation between 20 0/7 and 36 6/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*
Complication contributing to or causing death, choose **ONE** from the following list
- Infection
 - Meconium aspiration, other respiratory
 - Neurological issues/seizures
 - As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other
- b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)
Complication contributing to or causing death, choose **ONE** from the following list
- Infection
 - Meconium aspiration, other respiratory
 - Neurological issues/seizures

- As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other
- B. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if no continue, if yes revert to (A) above)
- a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*
- Complication contributing to or causing death, choose **ONE** from the following list
- Infection
 - Meconium aspiration, other respiratory
 - Neurological issues/seizures
 - As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other
- b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)
- Complication contributing to or causing death, choose **ONE** from the following list
- Infection
 - Meconium aspiration, other respiratory
 - Neurological issues/seizures
 - As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other

Neonatal Deaths (live born infant who died between day 7 and day 28 of extra-uterine life)

1. Did the death occur OOH (if yes continue, if no skip to (2) below)

A. County in which death diagnosed

B. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if yes continue, if no skip to (b) below)

a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a pregnancy or trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a pregnancy or trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

C. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if no continue, if yes revert to (A) above)

a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures

- As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - Other medical issue
 - Unknown
 - Information not available
 - Other
- b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

2. Did the death occur after transfer to another provider or facility? (if yes continue, if no revert to (1) above)

B. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if yes continue, if no skip to (b) below)

- a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

- b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

C. Was the death due to a diagnosed (either before or after delivery) anomaly deemed incompatible with life? (if no continue, if yes revert to (A) above)

- a. Was neonate born of a gestation between 20 0/7 and 36 0/7 weeks? (if yes continue, if no skip to (ii) below) *we could include a question asking the exact number of gestational weeks*

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

- b. Was neonate born of a gestation greater than 36 0/7 weeks? *we could include a question asking the exact number of gestational weeks* (if yes continue, if no revert to (a) above)

Complication contributing to or causing death, choose **ONE** from the following list

- Infection
- Meconium aspiration, other respiratory
- Neurological issues/seizures
- As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- Other medical issue
- Unknown
- Information not available
- Other

Maternal Deaths

1. Did the death occur OOH? (if yes continue, if no skip to (2) below)
 - A. County where death was diagnosed
 - B. Did the death occur prior to labor? (if yes continue, if no skip to (C) below)
 - a. Was gestation greater than 20 weeks (if yes skip to (D) below, if no continue)
 - b. Was gestation greater than 37 weeks (if yes skip to (D) below, if no continue)
 - c. Was gestation greater than 42 weeks (if yes skip to (D) below, if no revert to (a or b) above)
 - C. Complication contributing to or resulting in death. Mark all that apply:
 - i. Blood loss
 - ii. Sepsis
 - iii. Eclampsia/toxemia/HELLP syndrome
 - iv. Embolism (pulmonary or amniotic fluid)
 - v. As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - vi. As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - vii. Unknown
 - viii. Information not available
 - ix. Other
 - D. Did death occur during labor? (if yes continue, if no skip to (D) below)
 - a. Was gestation greater than 20 weeks (if yes skip to (D) below, if no continue)
 - b. Was gestation greater than 37 weeks (if yes skip to (D) below, if no continue)
 - c. Was gestation greater than 42 weeks (if yes skip to (D) below, if no revert to (a or b) above)
 - d. Complication contributing to or resulting in death. Mark all that apply:
 - i. Blood loss
 - ii. Sepsis
 - iii. Eclampsia/toxemia/HELLP syndrome
 - iv. Embolism (pulmonary or amniotic fluid)
 - v. As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - vi. As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - vii. Unknown
 - viii. Information not available
 - ix. Other
 - E. Did death occur after delivery and prior to the 42nd day post-partum? (if yes continue, if no revert to (B or C) above)
 - a. Was gestation greater than 20 weeks (if yes skip to (D) below, if no continue)
 - b. Was gestation 20 0/7 to 3 6/7 weeks (if yes skip to (D) below, if no continue)
 - c. Was gestation greater than 42 weeks (if yes skip to (D) below, if no revert to (a or b) above)
 - d. Complication contributing to or resulting in death. Mark all that apply:
 - i. Blood loss
 - ii. Sepsis

- iii. Eclampsia/toxemia/HELLP syndrome
- iv. Embolism (pulmonary or amniotic fluid)
- v. As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
- vi. As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
- vii. Unknown
- viii. Information not available
- ix. Other

2. Did the death occur after transfer to another provider or facility? (if yes continue, if no revert to (1) above)

A. County where death was diagnosed

B. Did the death occur prior to labor? (if yes continue, if no skip to (C) below)

- a. Was gestation greater than 20 weeks (if yes skip to (D) below, if no continue)
- b. Was gestation greater than 37 weeks (if yes skip to (D) below, if no continue)
- c. Was gestation greater than 42 weeks (if yes skip to (D) below, if no revert to (a or b) above)
- d. Complication contributing to or resulting in death. Mark all that apply:
 - i. Blood loss
 - ii. Sepsis
 - iii. Eclampsia/toxemia/HELLP syndrome
 - iv. Embolism (pulmonary or amniotic fluid)
 - v. As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - vi. As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - vii. Unknown
 - viii. Information not available
 - ix. Other

F. Did death occur during labor? (if yes continue, if no skip to (D) below)

- a. Was gestation greater than 20 weeks (if yes skip to (D) below, if no continue)
- b. Was gestation greater than 37 weeks (if yes skip to (D) below, if no continue)
- c. Was gestation greater than 42 weeks (if yes skip to (D) below, if no revert to (a or b) above)
- d. Complication contributing to or resulting in death. Mark all that apply:
 - i. Blood loss
 - ii. Sepsis
 - iii. Eclampsia/toxemia/HELLP syndrome
 - iv. Embolism (pulmonary or amniotic fluid)
 - v. As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - vi. As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - vii. Unknown
 - viii. Information not available
 - ix. Other

- G. Did death occur after delivery and prior to the 42nd day post-partum? (if yes continue, if no revert to (B or C) above)
- a. Was gestation greater than 20 weeks (if yes skip to (D) below, if no continue)
 - b. Was gestation greater than 37 weeks (if yes skip to (D) below, if no continue)
 - c. Was gestation greater than 42 weeks (if yes skip to (D) below, if no revert to (a or b) above)
 - d. Complication contributing to or resulting in death. Mark all that apply:
 - i. Blood loss
 - ii. Sepsis
 - iii. Eclampsia/toxemia/HELLP syndrome
 - iv. Embolism (pulmonary or amniotic fluid)
 - v. As a direct result of a trial of labor in a woman with a history of 1 or 2 prior cesarean deliveries
 - vi. As a direct result of a trial of labor in a woman with a history of more than 2 prior cesarean deliveries
 - vii. Unknown
 - viii. Information not available
 - ix. Other